

# Burnett County Native Plant Lists



## Moist Woodland Edge

Medium, loamy to wet, organic soils

Partial shade 4–8 hours

Common Name	Scientific Name	Height	Flower Color	Bloom Time	Butterflies	Hummingbirds	Songbirds	Waterfowl	Game Birds	Reptiles/Amphibian	Small Mammals	Deer
<b>Groundcover</b>												
Awl fruit sedge	<i>Carex stipata</i>	12-42"	NA	NA	•		•	•	•		•	•
Blue joint grass	<i>Calamagrostis canadensis</i>	3-5'	NA	NA			•	•			•	•
Caterpillar/Fringed sedge	<i>Carex crinita</i>	2-5'	NA	NA			•	•	•		•	•
Fox sedge	<i>Carex vulpinoidea</i>	18-30"	NA	NA	•		•	•	•		•	•
Fringed brome	<i>Bromus ciliatus</i>	2-4'	NA	NA			•	•	•		•	•
Prairie cordgrass	<i>Spartina pectinata</i>	4-8'	NA	NA			•		•		•	•
Virginia wild rye	<i>Elymus virginicus</i>	2-4'	NA	NA			•	•	•			•
<b>Shrubs</b>												
Am. mountain ash	<i>Sorbus americana</i>	to 28'	White	June			•		•			
Black chokeberry	<i>Aronia melanocarpa</i>	3-6'	White	May	•		•		•		•	•
Chokecherry°	<i>Prunus virginiana</i>	to 25'	White	May-June	•		•		•		•	•
Elderberry°	<i>Sambucus canadensis</i>	3-9'	White	June-July			•		•		•	•
Highbush cranberry	<i>Viburnum trilobum</i>	10-13'	White	June	•		•		•		•	•
Ironwood	<i>Carpinus caroliniana</i>	to 30'	NA	NA	•		•		•		•	•
Meadowsweet°	<i>Spiraea alba</i>	3-6'	White	June-Aug	•		•		•			•
Nannyberry°	<i>Viburnum lentago</i>	to 20'	White	June			•		•		•	
Northern bayberry	<i>Myrica gale</i>	3-4'	NA	NA			•		•			•
Pussy willow	<i>Salix discolor</i>	10-25'	White	April-May	•		•	•	•	•	•	•
Red osier dogwood°	<i>Cornus stolonifera</i>	4-10'	White	May-Aug	•		•	•	•		•	•
Sandbar willow	<i>Salix interior</i>	6-15'	White	April			•	•	•	•	•	•
Speckled alder	<i>Alnus incana</i>	10-25'	NA	NA			•	•	•		•	•
Steeplebush°	<i>Spirea tomentosa</i>	2-4'	Pink	July-Aug	•		•		•			•
Winterberry holly	<i>Ilex verticillata</i>	to 10'	White	June			•		•		•	•
<b>Trees</b>												
Black ash°	<i>Fraxinus nigra</i>	to 50'	NA	NA			•	•	•		•	•
Black willow°	<i>Salix nigra</i>	to 60'	NA	NA	•		•	•	•		•	•
Green ash°	<i>Fraxinus pensylvanica</i>	to 60'	NA	NA			•	•	•		•	•
Northern white cedar	<i>Thuja occidentalis</i>	to 70'	NA	NA			•				•	•
Paper birch	<i>Betula papyrifera</i>	to 70'	NA	NA			•		•		•	•
River birch	<i>Betula nigra</i>	to 75'	NA	NA			•	•	•		•	•
Silver maple°	<i>Acer saccharinum</i>	to 80'	NA	NA			•	•	•		•	•
Tamarack	<i>Larix laricina</i>	to 80'	NA	NA			•		•		•	•
White spruce°	<i>Picea glauca</i>	to 100'	NA	NA			•				•	•
Yellow birch	<i>Betula alleghaniensis</i>	to 80'	NA	NA			•	•	•	•	•	•

NA = Not Applicable - no flowers or inconspicuous flowers

° = Clay-tolerant plant

# Select Species



## Moist Woodland Edge

Medium, loamy to wet, organic soils

Partial shade 4–8 hours



Awl fruit sedge (*Carex stipata*)



Blue joint grass (*Calamagrostis canadensis*)



Fox sedge (*Carex vulpinoidea*)



Chokeberry (*Aronia melanocarpa*)



Chokecherry (*Prunus virginiana*)



High bush cranberry (*Viburnum trilobum*)



Meadowsweet (*Spiraea alba*)



Nannyberry (*Viburnum lentago*)



Red osier dogwood (*Cornus stolonifera*)



Steeplebush (*Spiraea tomentosa*)



Tamarack (*Salix nigra*)



Yellow birch (*Betula alleghaniensis*)